

# Hops: Diseases

Mizuho Nita, Extension Plant Pathologist,  
Alson H. Smith Jr. AREC

## ■ Nonchemical Approaches

### PREPLANTING CONSIDERATIONS

---

#### SITE SELECTION AND EVALUATION

Air circulation and water drainage are the two key factors when it comes to disease management. Poorly drained soil promotes some soilborne pathogens, such as black root rot (caused by *Phytophthora citricola*) and crown gall (caused by *Agrobacterium tumefaciens* aka *Rhizobium radiobacter*). Poor air circulation will promote diseases such as downy mildew and Botrytis gray mold because the pathogens for these diseases thrive in moist environments. Also, please note that variety selection can depend on site characteristics. For example, cultivars such as Magnum and Perle have been grown for centuries in European countries with well-drained and low pH (5.5-6.2) soil conditions. Therefore, it is recommended that you examine characteristics of your site, such as water drainage, pH, air circulation, sun exposure, availability of nutrients, etc., prior to the selection of cultivars.

#### CULTIVAR SELECTION

Both downy mildew (caused by *Pseudoperonospora humuli*) and powdery mildew (caused by *Podosphaera macularis*) are destructive diseases of hop. Considering environmental conditions during the growing season, it is best to select varieties that are not susceptible to these two diseases. Cascade, Fuggle, Magnum, Newport, and Perle are considered resistant to downy mildew. Comet, Crystal, First Gold, Newport, and Nugget are resistant to powdery mildew, and other cultivars, such as Cascade, Centennial, Hallertauer Tradition, Liberty, Pioneer, and Teamaker are moderately resistant to powdery mildew. On the other hand, cultivars such as Cluster, East Kent Golding, Tolhurst, and Vanguard are known to be susceptible to both powdery and downy mildew; thus, these cultivars should be avoided.

#### QUALITY OF RHIZOMES

The downy mildew pathogen as well as virus pathogens can survive in plant tissues. Therefore, it is very important to obtain certified rhizomes from reputable sources. Unfortunately, even certified, disease-tested rhizomes have a chance of carrying pathogens; however, certified rhizomes will be much cleaner than noncertified rhizomes and will greatly minimize the risk of disease development in young hop yards. The National Clean Plant Network has a program for hops since 2010, and producing 58 cultivars (<https://www.nationalcleanplantnetwork.org/hops>)

#### OTHER CULTURAL PRACTICES

Row spacing and row orientation need to be carefully planned, not only

to maximize production per acreage, but also to achieve good air circulation. For example, higher planting density with Cascade is known to increase the risk of downy and powdery mildew development. If irrigation is considered for your hop yards, overhead irrigation should be avoided because it will create an environment that is conducive to disease development. Also, excessive nitrogen can make hop plants more susceptible to some diseases.

### IN-SEASON CONSIDERATIONS

---

#### DORMANT SEASON DISEASE MANAGEMENT

Both downy mildew and powdery mildew pathogens can overwinter in infected dormant buds and crowns. The emerging buds can be infected during the winter and spring. Thus, it is important to manage downy mildew and powdery mildew as the season starts. Spring pruning can be done in late winter or early spring by removing all basal shoots to remove potentially infected shoots from the previous season. Also, “crowning,” the removal of the top 1-2 inches of the crown prior to training, and “scratching,” the removal of buds from within 1-2 inches of the soil surface using a special device, can be done. The timing of pruning is variety-specific and can affect yield potential; follow recommendations for your varieties.

#### STRIPPING

After the trained bines grow to a certain height, the lower 4-5 feet of the leaves and lateral branches need to be removed to minimize the spread of downy and powdery mildew. The process is called “stripping” and can be done mechanically.

## ■ Chemical Control Recommendations

### PATHOGEN BIOLOGY AND TIMING OF FUNGICIDE APPLICATION

---

Use of fungicides depends on other factors, such as site, weather, cultivar, cultural practice, etc. However, in general, preventative application of fungicides with 10-14-day intervals is recommended to minimize the risk of disease outbreak. The frequency of application depends on the growth of the bines as well as environmental conditions. For example, the downy mildew pathogen becomes active when the temperature is above 41 degrees Fahrenheit (F), and rain events promote their infection process. The powdery mildew pathogen becomes active at temperatures above 46°F, and the optimal temperature for pathogen growth is 64-70°F. The infection risk is very high when the minimum nighttime temperature is above 50°F and the daily high temperature is below 68°F. Once temperatures increase in the summer (3 or more hours above 86°F per day), the risk of powdery mildew infection decreases. Another disease to be considered is Botrytis gray mold, caused by *Botrytis cinerea*. When a prolonged wetness event is expected at burr development, a specific application against Botrytis may be needed.

## FUNGICIDE RESISTANCE MANAGEMENT

Use sufficient water to ensure complete coverage of the foliage. Make sure to (1) tank mix, (2) rotate the Fungicide Resistance Action Committee (FRAC) mode of action groups ([www.frac.info](http://www.frac.info)), and (3) limit the use of the same FRAC group to 2-3 times per season (with exceptions for copper, sulfur, oil, and other materials that has M or NC in their FRAC group) to minimize the risk of fungicide resistance development. Tank mixing of fungicide can be done by placing two or more fungicidal chemicals into a spray tank. It is recommend to mix two or more different mode of action groups, which target the same disease, in order to reduce development of fungicide resistance. The mode of action of a chemical is the way(s) for the chemical to either kill or deactivate the target pathogens, and its classification is listed on the label as FRAC group. Fungicides with the same FRAC group share the same mode of action and are essentially the same in terms of the risk of fungicide resistance. In order to minimize the cost of application, it is often recommended to mix a material with a

single mode of action and a broad-spectrum material, such as copper. Also, please note that some combinations of fungicides, oils, and plant nutrients are not compatible when mixed in the same tank, which may cause injury to your plant. Some labels list tank mixing partners.

In the case of a prolonged rain event, you may use either phosphorous acid (FRAC group P07) and/or metalaxyl (FRAC group 4) to control ongoing infection of downy mildew. However, application of the material has to be done within a few days of rain, prior to symptom development. Application of these materials against actively sporulating downy mildew colonies can increase the risk of fungicide resistance development. The same principle applies to fungicide resistance management of powdery mildew. Avoid the application of FRAC group 3, 7, 11, 13, 27, or U8 fungicide onto actively sporulating powdery mildew colonies. Potassium salt or oil-based products are recommended in such a case.

**Table 4.1 - Fungicides Registered for Control of Hop Diseases in Virginia**

Pest	Fungicide	Rate/Acre	FRAC* Grouping	Spray Timing and Remarks
<b>Before bine training</b>				
Downy mildew	Metastar 2E	1 qt	4	Timing and rate of application differ based on application method, row spacing, and product. Please refer to the label for the rate, amount of water, mixing partners, timing of application, etc.
	Ridomil Gold SL	0.5 pt	4	
	Ultra Flourish	1 qt	4	
	Orondis Gold	20.0-36.0 fl oz	4+49	
<b>From the beginning of bine training</b>				
Downy mildew	Curzate 60DF	3.2 oz	27	Ten to 14-day interval application (depends on the environmental conditions). See labels for mixing partners. Many modes of action (i.e., FRAC group) listed here are known for developing fungicide resistance. Always mix with a broad spectrum fungicide such as copper. For rainy season, consider mixing with a FRAC = P07 material.
	fixed copper (various formulations)	See label	M1	
	Flint Extra	See label	11	
	Forum	6.0 oz	40	With other crops, downy mildew strains that are resistant to Flint, Pristine, and other strobilurin (aka Qol) and Metastar (metalaxyl) fungicides are common. Rotating and tank mixing with a different anti-downy-mildew material (e.g., copper) is highly recommended.
	fosetyl-Al = Aluminum tris, various formulations	See label	P07	
	Luna Sensation	7.6 fl oz	7+11	
	Metastar 2E	1 qt	4	
	phosphorous acid (various formulations)	See label	P07	
	Orondis Gold	20.0-26.0 fl oz	4 + 49	
	Orondis Ultra	5.5-8.0 fl oz	40+49	
	Pristine	14.0 oz	7+11	
	Ranman 400SC	2.1-2.75 fl oz	21	
	Revus	8.0 oz	40	
	Ridomil Gold SL	0.5 pt	4	There are many formulations of phosphorous acid and fosetyl-Al (both are FRAC group P07) and copper products; refer to the label for the rate for the product. A high concentration of a phosphorous acid may cause plant injury. Also, some formulations of phosphorous acid and copper may cause plant injury (please refer to the label for more details).
	Tanos	8.0 oz	11+27	
	Ultra Flourish	1 pt	4	
Viathon	2.0 - 4.0 pt	P07 +3		
Zampro	11.0-14.0 fl oz	40 + 45		
				Group 40 materials (Forum, Revus, and Zampro) provide very good protection against downy mildew; however, as with others, tank mix, rotation, and limited usage (2-3 times/year) are recommended.
				Curzate, Metastar, or Tanos has to be tank mixed with another broad-spectrum fungicide active against downy mildew, such as copper.

Table 4.1 - Fungicides Registered for Control of Hop Diseases in Virginia (continued)				
Pest	Fungicide	Rate/Acre	FRAC* Grouping	Spray Timing and Remarks
Powdery mildew	fixed copper (various formulations)	See label	M1	10 to 14-day interval application (depends on the environmental conditions).
	Flint Extra	4.0 oz	11	Due to the risk of fungicide resistance development, do not use sterol inhibitors (aka DMI; FRAC group 3) or strobilurins (aka QoI; FRAC group 11) continuously; rotate with other groups of fungicides. It is recommended that sterol inhibitors and strobilurins be tank mixed with sulfur or copper material, and limit the use of FRAC groups 3 and 11 to 2-3 times/season. Pristine contains a strobilurin along with a different active chemical (SDHI) that has an efficacy against Botrytis gray mold.  Avoid using sterol inhibitors or strobilurins when there is a powdery mildew outbreak. Potassium bicarbonate products are recommended for an ongoing powdery mildew issue.  The rate for sulfur is generally 2-3 lb/100 gal, but it can be increased to as high as 6 lb/100 gal. Severe disease pressure may warrant this, but beware of possible plant injury at higher rates.  Mineral and neem oil products can control powdery mildew; however, there is some evidence of plant injury with the use of oil. See the label for detailed instructions. Also, avoid using oil and sulfur within two weeks of each other because it may cause injury to the plants.  If there is an outbreak of powdery mildew, consider using copper, oil, potassium bicarbonate, and sulfur.  If you plan to use Flint, Tanos or Ph-D, consider tank-mix with another FRAC material for fungicide resistance management.
	Luna Experience	8.0-17 fl oz	7+3	
	Luna Sensation	3.0-7.6 fl oz	7+11	
	mineral oil, neem oil	See label	NC	
	Ph-D	6.2 oz	19	
	potassium bicarbonate (Kaligreen, etc.)	See label	NC	
	Pristine	14.0 oz	7+11	
	triflumizole (Procure 480SC, etc.)	See label	3	
	Procure 480SC	4-8 lb	3	
	Quintec	4.0-8.2 oz	13	
	Rhyme	5.0-7.0 fl oz	3	
	Sonoma (various formulations)	See label	3	
	Tanos	8.0 oz	11+27	
	tebuconazole (various formulations)	2.0-8.0 oz	3	
	Torino	6.0-8.0 oz	U6	
	Trionic 4SC	12 fl oz	3	
Velum Prime	6.5-6.84 fl oz	7		
Viathon	2.0 - 4.0 pt	P07 +3		
Vivando	15.4 fl oz	U8		
wettable sulfur (various formulations)	2.0-6.0 lb	M2		
<b>After burr development</b>				
Botrytis	Pristine	14.0 oz	7+11	Although Pristine contains two different FRAC groups, strains of Botrytis that can overcome both FRAC groups are present in Virginia. Make sure to tank mix with a broad-spectrum fungicide, such as copper. Also, applications of Pristine should be limited to less than three times a season. Many biological control agents (e.g., Double Nickel, Howler, Serenade, etc.) list Botrytis on their labels. If you decided to use them, make sure to apply before potential infection event (i.e., rain). Also it is probably a good idea to test it in a small area before you apply broadly to make sure it fits your growing conditions.
	Luna Experience	8.0-17.0 fl oz	7+3	
	Luna Sensation	3.0-7.6 fl oz	7+11	
<b>After harvest</b>				
Downy mildew and powdery mildew	fixed copper (various formulation)	See label	M1	Once or twice after harvest. It is important to keep bines healthy after harvest to ensure the accumulation of carbohydrates into the rhizome for winter survival. Copper products should be sufficient to provide protection against both downy mildew and powdery mildew in most cases, but you may need to use sulfur to control powdery mildew. If there is a prolonged rain event and your hop yards have a history of downy mildew, you may need to use either phosphorous acid or fasetyl-Al.
	fosetyl-Al = Aluminum tris, various formulations	See label	P07	
	phosphorous acid (various formulations)	See label	P07	
	wettable sulfur (various formulations)	2-6 lb	M2	

\* Fungicide Resistance Action Committee.

<b>Table 4.2 - Product and Chemical Names, Restricted Entry Interval (REI), and Preharvest Interval (PHI)</b>			
<b>Product (chemical name) or if there are multiple products of the same chemical, Chemical name [product name]</b>	<b>Manufacturer</b>	<b>REI (hours)</b>	<b>PHI (days)</b>
Actinovate AG ( <i>Streptomyces lydicus</i> WYEC 108)	Novozimes BioAg Inc	4	0
Aliette WDG (Aluminum tris)	Bayer CropScience	4	0
Bio-Tam 2.0 ( <i>Trichoderma gamsii</i> , strain ICC 080, and <i>Trichoderma asperellum</i> , strain ICC 012)	SePRO Corp	4	0
Carbon Defense (potassium silicate)	FBSciences	4	0
Curzate 60DF (cymoxanil)	Corteva	12	7
Double Nickel 55 and LC ( <i>Bacillus amyloliquefaciens</i> , strain D747)	Certis USA	4	0
fixed copper [various formulations: Champ WG, Cueva, Kocide, Nordox 75WG, etc.]	Various	4-48 (see label)	0-14 (see label)
Empire (pyraclostrobin)	Sharda USA	12	14
Flint Extra (trifloxystrobin)	Bayer CropScience	12	14
Forum (dimethomorph)	BASF	12	7
fosetyl-Al [aluminum tris; Aliette WDG, Linebacker WDG]	Various	12	24
Gatten (flutianil)	Nichino America	12	7
Howler and Howler EVO ( <i>Pseudomonas chlororaphis</i> strain AFS009)	AgBiome	4	0
hydrogen dioxide or peroxide [OxiDate 5.0, Jet-Ag, etc.]	Various	4 (see label)	0 (see label)
JMS Stylet-Oil (paraffinic oil)	JMS Flower Farms	4	0
Luna Experience (fluopyram + tebuconazole)	Bayer	12	14
Luna Sensation (fluopyram + trifloxystrobin)	Bayer	12	14
metalaxyl [MetaStar 2E, ReCon 4F, etc.]	LG Life Sciences	48	45
mineral oil [various formulations: Damoil, Omni oil, Suffoil-X, etc.]	Various	4	0
neem oil [various formulations: Trilogy, Green Light, etc.]	Various	4	0
Orondis Gold (Oxathiapiprolin)	Syngenta	48	45
Orondis Ultra (Oxathiapiprolin)	Syngenta	4	7
Oxidate 2.0 or 5.0 (hydrogen peroxide and peroxyacetic acid)	BioSafe Systems	1	0
OxiPhos (phosphorous acid plus hydrogen peroxide)	BioSafe Systems	4	0
PerCarb (sodium carbonate peroxyhydrate)	BioSafe Systems	0	0
Ph-D (polyoxin D zinc salt)	UPL	4	0
phosphorous acid [phosphite; various formulations: Agri-Fos, Prophyt, Phostrol, etc.]	Various	4	0
potassium bicarbonate [various formulations: Armicarb, Kaligreen, etc.]	Various	4	0-1 (see label)
Pageant Intrinsic (boscalid)	BASF	12	14
Presidio (fluopicolide)	Valent	12	24
Pristine (pyraclostrobin + boscalid)	BASF	12	14
Procure 480SC	UPL	12	7
Quintec (quinoxifen)	Dow Agrosciences, GOWAN	12	21
myclobutanil [Rally, Sonoma, etc]	Various	24	14
Ranman (cyazofamid) (various cyazofamid formulations available)	FMC, Summit Agro	12	3

Regalia, Regalia PTO, and Regalia Rx ( <i>Reynoutria sachalinensis</i> )	Marrone Bioinnovations, Pro Farm Group	4	0
Revus (mandipropamid)	Syngenta	4	7
Rhyme (flutriafol)	FMC	12	7
Ridomil Gold SL (Mefenoxam) (various mefenoxam formulations available)	Syngenta	48	45
Serenade (various formulations)	Bayer Cropscience	4	0
Sodium Tetraborohydrate Decahydrate (two PREV-AM formulations)	Oro Agri	12	0
Sonata (various formulations)	Bayer Cropscience	4	0
sulfur [various formulations: Acoidal, Kumulus DF, Microthiol D, etc.]	Various	24	0
Tanos (cymoxanil + famoxadone)	Dupont, Corteva	12	7
tebuconazole [various formulations: AmTide Tebu 3.6 F, Orius, etc.]	Various	12	14
Tetraborohydrate Decahydrate (two PREV-AM formulations)	Oro Agri	12	0
triflumizole [various formulations: Procure, Trionic, Arkos, etc.]	Various	12 (see label)	14 (see label)
TRIONIC 4SC (Triflumizole)	UPI	12	7
Ultra Flourish (mefenoxam)	Nufarm	48	45
Velum Prime	Bayer CropScience	12	7
Viathon (phosphorous acid plus tebuconazole)	Helena	12	14
Vivando <sup>2</sup> (metrafenone)	BASF	12	3
Xyler FC (metalaxyl)	Vive Crop Protection, Inc	48	45
Zampro (amedoctradin + dimethomorph)	BASF	12	7
Zayin (geraniol)	GroPro Corporation	0	0
ZONIX (rhamnolipid)	Sepan Company	4	0
<sup>1</sup> Label recommends termination of use at burr development. <sup>2</sup> An older label does not list hops. A supplemental label is available.			



# Hops: Insects

*Douglas G. Pfeiffer, Fruit Entomologist, Virginia Tech*

## ■ Nonchemical Approaches

### SITE SELECTION

The likelihood that some insects will develop into pest status in a commercial planting is affected by environmental factors. This is true for the Japanese beetle. Larvae of the Japanese beetle feed on grass roots; consequently, if a planting is adjacent to pastureland, there is a nearby breeding ground. Plantings surrounded by woods will have less immigration pressure. Twospotted spider mite is favored by hot temperatures.

### BIOLOGICAL CONTROL

Some pests, such as the twospotted spider mite, are excellent candidates for biological control. A complex of native predators will feed on the twospotted spider mite and can maintain its populations

below damaging levels unless disrupted by chemical controls for other pests. Toxicity to natural enemies should be considered when deciding on a chemical control program. If disruptive materials are to be used, pay close attention to twospotted spider mite populations.

## ■ Chemical Control Recommendations

### RESISTANCE MANAGEMENT

Several factors affect the likelihood of resistance to pesticides, notably the number of generations per season and the number of offspring per female (fecundity). Spider mites are notorious for developing resistance to acaricides because they have a large number of generations annually and they produce many eggs. It is important to (1) rely on natural controls as much as possible and (2) obtain maximum coverage and rotate insecticides of different modes of action when chemical control is needed. Modes of action are indicated by the Insecticides Resistance Action Committee (IRAC) grouping (<https://irac-online.org/documents/moa-classification/?ext=pdf>).

Pest	Insecticide	Rate/Acre (unless noted)	IRAC* Grouping	Spray Timing and Remarks
Variegated cutworm, loopers	Baythroid XL	3.2 fl oz	3A	An action threshold has not been established for these pests. Larvae may cause defoliation on the plant and may feed on cones directly, especially late in the season. Larval population should not be allowed in the upper canopy in the late season. Sample by spreading a sheet on the ground and shaking the bine vigorously for about 15 seconds.  Dipel, Entrust and Venerate are OMRI-approved, suitable for organic production.
	Brigade 2SC	3.8-6.4 fl oz	3A	
	Coragen 1.67EC	3.5-5 fl oz	28	
	Delegate 25WG	2.5-4.0 oz	5	
	Dipel DF	1 lb	11A	
	Entrust 2SC	4.0-6.0 fl oz	5	
	Venerate	4.0-8.0 qt	—	
Twospotted spider mite	Acramite 50WS	0.75-1.5 lb	UN	Do not spray for spider mites preventatively in order to prevent the development of resistance. If multiple sprays are needed, rotate to a different IRAC class. Savey is strictly an ovicide and will not immediately reduce a high population. An action threshold has not been established, but a tentative threshold is 2-3 female mites/leaf until mid-July and 10 mites/leaf thereafter. Use a 10x hand lens when counting mites. Natural enemies often control spider mites, so sprays should not be used against sub-economic populations. Use of pyrethroid insecticides will often induce mite outbreaks. Acramite may be used only once per season.
	Agri-Mek 8SC	1.75-3.5 fl oz	6	
	Envidor 2SC	18.0-24.7 fl oz	23	
	Portal 0.4E	2.0-3.0 pt	21A	
	Savey 50DF	4.0-6.0 oz	10A	
	Zeal	3.0-4.0 oz	10B	
	Kanemite	31 fl oz	20B	
Aphids	Admire Pro (foliar)	2.8 fl oz	4A	Before flowering, use a tentative action threshold of 5-10 aphids/leaf. Control is difficult if aphids become established on cones. BotaniGard is OMRI-approved, suitable for organic production. Venerate provides only suppression for aphids, and is OMRI-approved. Aza-Direct is also OMRI-approved.
	Admire Pro (soil)	2.8-8.4 fl oz		
	Aza-Direct	1-2 pt		
	Baythroid XL	3.2 fl oz	3A	
	Beleaf 50SG	1.7-2.8 oz		
	BotaniGard ES	1 qt	N/A	
	Brigade 2SC	3.8-6.4 fl oz	3A	
	Fulfill 50WDG	4.0-6.0 oz	9B	
	Malathion 57EC	1 pt	1B	
	Movento 2EC	5.0-6.0 fl oz	23	
	Sivanto	7.0-10.5 fl oz	4D	
	Venerate	4-8 qts	—	

Pest	Insecticide	Rate/Acre (unless noted)	IRAC* Grouping	Spray Timing and Remarks
European corn borer	Baythroid XL	3.2 fl oz	3A	
	Brigade 2SC	3.8-6.4 fl oz	3A	
	Dipel DF	1 lb	11A	
	Coragen 1.67EC	3.5-5.0 fl oz	2B	
Japanese beetle	Aza-Direct	1-2 pt		Surround, Neemix, Trilogy, Aza-Direct, and PyGanic are all OMRI-approved. Neemix and Trilogy are to be applied together. Aza-Direct, Neemix and Trilogy should be applied before JB reaches high levels.
	Brigade 2SC	3.8-6.4 fl oz	3A	
	Neemix 4.5 + Trilogy	1-2 gal Neemix + 7.0-16.0 fl oz Trilogy/100 gal	UN	
	PyGanic 5EC	16-32 fl oz/100 gal	3A	
	Surround	25-50 lb	UN	
Potato leafhopper	Malathion 57EC	1 pt	1B	Potato leafhopper may be more common if alfalfa is grown nearby, especially after alfalfa cutting. M-Pede is OMRI-approved.
	M-Pede	2% solution	—	
	Movento 2SC	5-6 fl oz	23	
Question mark	Aza-Direct	1-2 pt		
	Delegate 25WG	2.5-4 oz	5	
	Botanigard Maxx	2 qt	3A	
	Dipel DF	1 lb	11A	
	Coragen 1.67EC	3.5-5.0 fl oz	2B	
Spotted lanternfly	Baythroid XL	3.2 fl oz	3A	Spotted lanternfly is a new invasive pest in the state. SLF was initially located in northern Virginia, but has been expanding its geographic range significantly. While this insect feeds on more than 70 different host plants, hop may be a host plant. Feeding by both nymphs and adults produces a large amount of honeydew, which may contaminate cones. Check cones for contamination if SLF is present on bines. Please read the SLF material posted at <a href="https://www.virginiafruit.ento.vt.edu/SLF.html">https://www.virginiafruit.ento.vt.edu/SLF.html</a> . This site has identification and control information, as well as updated distribution maps.
	Brigade 2SC	3.8-6.4 fl oz	3A	
Stink bugs	Beleaf 50SG	1.0-2.8 oz		

\*Insecticides Resistance Action Committee

Pesticide	Manufacturer	REI (hours)	PHI (days except where noted)
Acramite (bifenazate)	Chemtura	12	14
Admire Pro	Bayer	12	60 (soil), 28 (foliar)
Agri-Mek (abamectin)	Syngenta	96	28
Aza-Direct (azadirachtin)	Gowan	4	0
Baythroid (beta-cyfluthrin)	Bayer	12	7
Beleaf (flonicamid)	FMC	12	10
BotaniGard ES ( <i>Beauveria bassiana</i> )	LAM International	4	0
Botanigard Maxx ( <i>Beauveria bassiana</i> )	LAM International	12	0
Brigade (bifenthrin)	FMC	12	14
Coragen (chlorantraniliprole)	Dupont	4	0
Delegate (spinetoram)	Dow	4	1
Dipel ( <i>Bacillus thuringiensis</i> )	Valent	4	0
Entrust SC (spinosad)	Dow	4	1
Envidor (spirodiclofen)	Bayer	12	14
Fulfill (pymetrozine)	Syngenta	12	14



Table 4.4 - Insecticide Names, Restricted Entry Interval (REI), and Preharvest Interval (PHI) (continued)			
Pesticide	Manufacturer	REI (hours)	PHI (days except where noted)
Kanemite (acequinocyl)	Arysta	12	7
Malathion (malathion)	Loveland	12	10
Movento (spirotetramat)	Bayer	24	7
M-Pede (insecticidal soap)	Gowan	12	0
Neemix (azadirachtin)	Certis	4	0
Portal (fenpyroximate)	Nichino	12	15
PyGanic (pyrethrins)	MGK	12	0
Savey 50DF (hexythiazox)	Gowan	12	Until burr formation
Sivanto Prime (flupyradifurone)	Bayer	4	21
Surround (kaolin)	NovaSource	4	0
Trilogy (clarified extract of neem oil)	Certis	4	0
Venerate (Burkholderia)	Marrone	4	0
Zeal (etoxazole)	Valent	12	7
<b>A new invasive insect</b>			
<p>In January 2018, a new invasive insect was found in Virginia. Spotted lanternfly came to Virginia from southeastern Pennsylvania, and had been expanding its range within Frederick County. SLF feeds on more than 70 different hosts, and can cause significant injury on some. Some of our important fruit crops are on the host list: grape, caneberry, blueberry, stone and pome fruits, and hops. There have been conflicting reports on severity of injury to hops in Pennsylvania; for the time being, hop growers are urged to be vigilant. Populations can build to create a severe nuisance in residential areas as well. An eradication effort has been implemented in 2018, and a quarantine was established by VDACS in May 2019. More information on the quarantine program is posted (<a href="https://www.pubs.ext.vt.edu/ENTO/ENTO-319/ENTO-319.html">https://www.pubs.ext.vt.edu/ENTO/ENTO-319/ENTO-319.html</a>). An online training is available too allow certification as part of the quarantine effort (<a href="https://register.ext.vt.edu/search/publicCourseSearchDetails.do?method=load&amp;courseId=210837&amp;selectedProgramAreaId=25577&amp;selectedProgramStreamId=">https://register.ext.vt.edu/search/publicCourseSearchDetails.do?method=load&amp;courseId=210837&amp;selectedProgramAreaId=25577&amp;selectedProgramStreamId=</a>). So far, SLF has not caused economic loss in our agricultural crops. Nevertheless, hop growers should be aware of the pest's appearance, and how to handle finds you may make in your operations. For updated information, visit the spotted lanternfly page in the Virginia Cooperative Extension web site (<a href="https://ext.vt.edu/">https://ext.vt.edu/</a>). For updated control information, visit the SLF page in Virginia Fruit (<a href="https://www.virginiafruit.ento.vt.edu/SLF.html">https://www.virginiafruit.ento.vt.edu/SLF.html</a>). To report suspected discoveries, please visit the SLF reporting page (<a href="https://ask.extension.org/groups/1981/ask">https://ask.extension.org/groups/1981/ask</a>).</p>			



# Hops: Weed Management

*Jeffrey Derr, Extension Weed Scientist, Hampton Roads AREC*

## ■ Nonchemical Approaches

Apply mulch at a 2-4 inch depth using a mulch that is free of weed seed or use shallow cultivation (2-4 inches).

## ■ Chemical Control Recommendations

### ORGANIC PRODUCTION

Acetic acid (Weed Pharm) can be applied as a directed spray for nonselective contact control of small annual weeds. This will suppress perennial weeds; repeat treatments will be needed to control regrowth. Do not allow spray to contact hops foliage.

### CONVENTIONAL PRODUCTION

Application	Common Name (Trade Name)	Rate of Active Ingredient per Acre (Rate of Product per Acre)	Timing and Remarks
Preemergence to weeds	flumioxazin (Chateau)	3 oz (6 oz)	Apply to dormant hops in January-March as a 1-1.5-foot strip on each side of the row. Make only 1 application/year. For small areas, apply 0.13 oz Chateau/1,000 sq ft. Controls a range of annual weeds
	pendimethalin (Prowl H2O)	1.0-4.0 lb (1.1-4.2 qt/A)	Do not apply otop hops; instead apply as a direct spray. Do not apply more than 4.2 qt/A/year. For small areas apply 0.8-3.0 fl oz/1000 sq ft. PHI is 90 days.
	norflurazon (Solicam)	2-4 lb (2.5-5.0 lb)	Apply as a directed spray to hops established at least 6 months. Use lower rates on sandy soils. For small areas, apply 0.9-1.8 oz Solicam/1,000 sq ft. Controls a range of annual weeds and suppresses yellow nutsedge.
	trifluralin (Treflan 4EC or other labeled formulation)	0.50-0.75 lb (1.0-1.5 pt)	Apply as a directed spray to dormant, established hops, avoiding hops crowns. Immediately incorporate 1-2 inches deep. Do not spray over hops. For small areas, apply 0.37-0.55 fl oz Treflan 4EC/1,000 sq ft. Controls annual grasses and certain small-seeded annual broadleaf weeds.
Postemergence to weeds	carfentrazone (Aim EC)	0.016-0.031 lb (1.0-2.0 fl oz/A for broadleaf weeds, 2.0 fl oz/A for sucker management)	Contact herbicide for control of broadleaf weeds less than 4 inches tall using a hooded sprayer. Can also be applied post-directed for sucker management. Add a nonionic surfactant or crop oil concentrate. Apply after trained hops stems are woody when used for sucker management.
	clethodim (Select Max)	0.07-0.12 lb (9-16 fl oz)	Apply to actively growing annual and perennial grassy weeds. Will not control nongrass monocots or any broadleaf weeds. Perennial grasses like johnsongrass and bermudagrass generally will require retreatment. Add a nonionic surfactant at 0.25% V/V. There is a 21-day PHI.
	2,4-D (2,4-D Amine 4 or other labeled formulation)	0.48 lb ae (1 pt)	Make a directed spray to the row middles. Use a shielded or hooded sprayer to prevent contact with hops foliage. Controls certain annual broadleaf weeds. The PHI is 28 days.
	glyphosate (Roundup PowerMAX or other labeled formulation)	0.39-3.70 lb ae (11.0 fl oz-3.3 qt)	Apply only when there are no green shoots of hops within the spray zone. For small areas, mix 2 fl oz/gallon and lightly wet the weed foliage. There is a 14-day PHI. Controls annual and perennial weeds. Use a shielded spray to avoid contact with hops foliage.
	pelargonic acid (Scythe)	(3-7% V/V)	Rapid acting contact herbicide. Treat weeds when they are less than 4 inches tall. Avoid contact with hops foliage except when used for sucker management. Apply after trained hops stems are woody when used for sucker management.

<b>Table 4.6 - Herbicide, Manufacturer, Restricted Entry Interval (REI), and Preharvest Interval (PHI)</b>			
<b>Herbicide Trade Name</b>	<b>Manufacturer</b>	<b>REI (hours)</b>	<b>PHI (days)</b>
2,4-D Amine 4	WinField Solutions	48	28
Aim	FMC	12	7
Chateau	Valent	12	30
Prowl H2O	BASF	24	90
Roundup PowerMax	Bayer	4	14
Scythe	Gowan	12	1
Select Max	Valent	24	21
Solicam	Tessenderlo Kerley	12	60
Treflan 4EC	Helena	12	—

